Exam #2 · Thursday, March 1, 2007

MATH 124 · Calculus I · Section 8 · Spring 2007	Name
Problem 1. Let $f(x) = x^x$. Numerically approximate three successively smaller values of h .	f'(2) using difference quotients. Use at least
Problem 2. On a mountain-climbing expedition, you find y be your altitude above sea level, measured in feet; let H	
Part (a). What are the units of $H'(y)$?	
Part (b). What is the sign of $H'(y)$?	
Part (c). Give a practical interpretation of $H^{-1}(35)$.	
Part (d). Give a practical interpretation of $H'(7500)$.	
Problem 3. In a lab experiment, you have microorgan of microganisms, in millions, is a function of time t in number is given by	

 $m(t)=4.1e^{0.24t}.$ Find the rate of change in population on day 5 of the experiment. In your answer, please show units.

Problem 4. Let $G(t) = 2^{-rt} \sin(at)$. Find G'(t).


Problem 5. Let $f(z) = \tan^{-1}(z)$. Compute f''(1).

Problem 6. Let $q(x) = \ln(2 + 2x + x^2)$.

Part (a). Find q'(x).

Part (b). Find an equation for the tangent line to q(x) at x=3.

Problem 7. Let F(t) and G(t) be given by the following graphs.



Part (a). Find G'(40).

Part (b). Let $H(t) = \frac{F(t)}{G(t)}$. Find H'(50).

Part (c). Let H(t) = F(G(t)). Find H'(30).

Problem 8. Let

$$f(x) = \begin{cases} \sin(x), & x \ge 0\\ x - x^3, & x \le 0. \end{cases}$$

Part (a). Find f'(x). Write it as a piecewise function.

Part (b). Is the original function f(x) differentiable at x = 0? Why or why not? (Hint: graph it.)

Problem 9. When is $g(x) = x^3 + bx^2 + cx + d$ concave up? Assume b, c, d are constants. (You will need to solve an inequality.)