

Numerical methods for random spatial permutations

Stochastic Processes and Their Applications, Berlin, 2009

John Kerl

Department of Mathematics, University of Arizona

July 30, 2009

The probability model


State space: $\Omega_{\Lambda, N} = \Lambda^N \times \mathcal{S}_N$, where $\Lambda = [0, L]^3$ with periodic boundary conditions.

Point positions: $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_N)$ for $\mathbf{x}_1, \dots, \mathbf{x}_N \in \Lambda$.

Hamiltonian, where $T = 1/\beta$ and $r_\ell(\pi)$ is the number of ℓ -cycles in π :

$$H(\mathbf{X}, \pi) = \frac{T}{4} \sum_{i=1}^N \|\mathbf{x}_i - \mathbf{x}_{\pi(i)}\|^2 + \sum_{\ell=1}^N \alpha_\ell r_\ell(\pi).$$

- The **first term** discourages long permutation jumps, moreso for higher T .
- The **temperature** scale factor $T/4$, not $\beta/4$, is surprising but correct for the Bose-gas derivation of the Hamiltonian.
- The **second term** discourages cycles of length ℓ , moreso for higher α_ℓ . These **interactions** are not between points, but rather between **permutation jumps**.



The probability model


Fixed point positions (**quenched model** — includes all simulations done up to the present on the **lattice** $N = L^3$):

$$P_{\mathbf{X}}(\pi) = \frac{1}{Y(\Lambda, \mathbf{X})} e^{-H(\mathbf{X}, \pi)}, \quad Y(\Lambda, \mathbf{X}) = \sum_{\sigma \in \mathcal{S}_N} e^{-H(\mathbf{X}, \sigma)}.$$

Varying positions (**annealed model** — many theoretical results are available):

$$P(\pi) = \frac{1}{Z(\Lambda, N)} e^{-H(\mathbf{X}, \pi)}, \quad Z(\Lambda, N) = \frac{1}{N!} \int_{\Lambda^N} Y(\Lambda, \mathbf{X}) d\mathbf{X}.$$

In either case, we write the **expectation** of an RV as $\mathbb{E}_{\pi}[\theta(\pi)] = \sum_{\pi \in \mathcal{S}_N} P(\pi)\theta(\pi)$.



The probability model: intuition



What does a random spatial permutation actually look like? (Recall

$$H(\mathbf{X}, \pi) = \frac{T}{4} \sum_{i=1}^N \|\mathbf{x}_i - \mathbf{x}_{\pi(i)}\|^2 + \sum_{\ell=1}^N \alpha_{\ell} r_{\ell}(\pi).)$$

- As $T \rightarrow \infty$, the probability measure becomes supported only on the **identity permutation**. For large but finite T : there are tiny islands of 2-cycles, 3-cycles, etc.
- As $T \rightarrow 0$, length-dependent terms go to zero. The probability measure approaches the **uniform distribution** on \mathcal{S}_N : all π 's are equally likely.

For intermediate T , things get more interesting:

- The length of each permutation jump, $\|\pi(\mathbf{x}) - \mathbf{x}\|$, remains small.
- For T above a **critical temperature** T_c , all cycles are short: 2-cycles, 3-cycles, etc. $T_c \approx 6.8$, and positive α terms increase T_c .
- **Phase transition** at T_c : for $T < T_c$ jump lengths remain short but *long cycles form*.
- Figures: high T , medium but subcritical T , and low T .



Quantifying the onset of long cycles

We observe the following:





- For $T > T_c$, $\mathbb{E}[\ell_{\max}]$ is constant as $N \rightarrow \infty$: cycles remain **finite**.
- For $T < T_c$, $\mathbb{E}[\ell_{\max}]$ scales with N : there are arbitrarily long cycles, or **infinite cycles**, in the infinite-volume limit. **Feynman (1953)** studied long cycles in the context of Bose-Einstein condensation for interacting systems. See also **Sütő (1993, 2002)**.

Other random variables (“order parameters”) besides $\mathbb{E}[\ell_{\max}/N]$:

- **Fraction of sites in long cycles**, f_I , goes to zero in L above T_c , non-zero below.
- **Correlation lengths** $\xi(T)$ which are (spatial or hop-count) length of the cycle containing the origin: for $T < T_c$, these blow up in L .
- **Winding numbers**: number of x, y, z wraps around the 3-torus (Λ with p.b.c.). Scaled winding number: $f_S = \frac{\langle \mathbf{W}^2 \rangle L^2}{3\beta N}$. This behaves much like f_I , but is easier to compute with. Also, f_W : fraction of sites which participate in winding cycles.

Central goal of my dissertation work: quantify the **dependence of T_c on α** , where $\Delta T_c(\alpha) = \frac{T_c(\alpha) - T_c(0)}{T_c(0)}$. Known results and conjectures are formulated quantitatively in terms of $\lim_{\alpha \rightarrow 0} \Delta T_c(\alpha)$.

Behavior of order parameters as functions of L and T ($\alpha_\ell \equiv 0$)



Known results and conjectures

Recall $H(\mathbf{X}, \pi) = \frac{T}{4} \sum_{i=1}^N \|\mathbf{x}_i - \mathbf{x}_{\pi(i)}\|^2 + \sum_{\ell=1}^N \alpha_\ell r_\ell(\pi)$. We have the following models:

- **Non-interacting model:** $\alpha_\ell \equiv 0$.
- **Two-cycle model:** $\alpha_2 = \alpha$ and other cycle weights are zero.
- **Ewens model:** α_ℓ is constant in ℓ .
- **General-cycle model:** No restrictions on α_ℓ .

Known results for the continuum (obtained largely using Fourier methods):

- $\Delta T_c(\alpha)$ is known (to **first order in α**) for two-cycle interactions (Betz and Ueltschi, CMP 2008) and **small cycle weights** (Betz and Ueltschi 2008). (This taps into a long and controversial history in the physics literature: see Baym et al., EJP B 2001, or Seiringer and Ueltschi, PRB 2009, for surveys.) The critical (ρ, T, α) manifold relates ρ_c to T_c .

$$\rho_c(\alpha) \approx \sum_{\ell \geq 1} e^{-\alpha_\ell} \int_{\mathbb{R}^3} e^{-\ell 4\pi^2 \beta \|\mathbf{k}\|^2} d\mathbf{k} = \frac{1}{(4\pi\beta)^{3/2}} \sum_{\ell \geq 1} e^{-\alpha_\ell} \ell^{-3/2}$$

$$\Delta T_c(\alpha) \approx c\rho^{1/3}\alpha, \quad \text{for } \alpha \approx 0.$$

Known results and conjectures

Known results (continued):

- $\langle \ell_{\max} \rangle / N f_I$ is constant for $T < T_c$ for $\alpha_\ell \equiv 0$. (That is, the two order parameters f_I and $\langle \ell_{\max} \rangle / N$ have the same critical exponent.) For **uniform-random permutations** (Shepp and Lloyd 1966 solved Golomb 1964), $\langle \ell_{\max} \rangle / N \approx 0.6243$; unpublished work of Betz and Ueltschi has found $\langle \ell_{\max} \rangle / N f_I$ is that same number for the **non-interacting case** $\alpha_\ell \equiv 0$. Intuition: long cycles are “uniformly distributed” within the zero Fourier mode.

Conjectures:

- $\langle \ell_{\max} \rangle / N f_I$ is constant for $T < T_c$ for all interaction models. Questions: Why should this be true on the lattice? How does that constant depend on α ?
- $\xi(T)$ is monotone in T : currently unproved either for the continuum or the lattice.
- $\rho_c(\alpha)$ formula holds not only for **small cycle weights** ($\alpha_\ell \rightarrow 0$ faster than $1/\log \ell$).

Open questions:

- To what extent does the $\rho_c(\alpha)$ formula hold true on the **lattice**?
- $\Delta T_c(\alpha)$ on the **lattice** should be similar to that on the continuum.
- $\Delta T_c(\alpha)$ is theoretically unknown for **Ewens interactions** (continuum or lattice).

Metropolis sampling

The **expectation** of a random variable θ (e.g. ℓ_{\max}/N , f_I , f_S , ξ) is

$$\mathbb{E}_{\pi}[\theta(\pi)] = \sum_{\pi \in \mathcal{S}_N} P(\pi)\theta(\pi).$$

The number of permutations, $N!$, grows intractably in N . The expectation is instead **estimated** by summing over some number M (10^4 to 10^6) typical permutations.

The usual technical issues of Markov chain Monte Carlo (MCMC) methods are known and handled in my simulations and dissertation: **thermalization** time, proofs of **detailed balance**, **autocorrelation**, **batched means**, and **quantification of variance** of samples.


Metropolis step (analogue of single spin-flips for the Ising model): swap permutation arrows which end at nearest-neighbor lattice sites. This either splits a common cycle, or merges disjoint cycles:



As usual, the **proposed** change is **accepted** with probability $\min\{1, e^{-\Delta H}\}$.

Metropolis sampling and winding numbers: the GKU algorithm

- Figure part 1: a **long cycle** on the torus almost meets itself in the x direction.
- Part 2: after a Metropolis step, one cycle winds by $+1$, and the other by -1 . Metropolis steps create winding cycles only in **opposite-direction pairs**; total $W_x(\pi)$ is still zero.
- Part 3: if we **reverse one cycle** (zero-energy move), $W_x(\pi)$ is now 2.





Our current best algorithm (GKU) has two types of sweeps: (1) For each lattice site, do a **Metropolis step** as above (Gandolfo, K). (2) For each cycle in the permutation, **reverse** the direction of the cycle with probability $1/2$ (Ueltschi). This permits winding numbers of even parity in each of the three axes.

Methods for obtaining winding numbers of all parities: try (so far with mixed success) to adapt **non-local updates** (e.g. Swendsen-Wang for Ising) and **worm algorithm**. Problems with **low acceptance rate** and **stopping time for worm closure**, respectively.

Computational results: ΔT_c




These results are preliminary. For fixed L , one may sandwich $T_c(L)$ between the vertical asymptotes of $1/f_S$ and ξ . From such graphs, we obtain, for $L = 40$, with points on the lattice,

- $\Delta T_c(L)/\alpha = 0.0759 \pm 15\%$ for the r_2 model (vs. 0.088 theoretically for the continuum), and
- $\Delta T_c(L)/\alpha = 0.483 \pm 10\%$ for the Ewens model (theoretical value is unknown, but small-cycle-weight prediction for the continuum is 0.66).




Computational results: GRU quotient $\langle \ell_{\max} \rangle / N f_I$

The GRU quotient varies with α in the Ewens model, but not in the r_2 model. For small L , it is non-constant for $T < T_c$; this bias seems to disappear as $L \rightarrow \infty$. (Needs a statistical confidence test.)



For r_2 , GRU quotient is ≈ 0.626 regardless of α . For Ewens, averaging at all subcritical T 's, we get the following dependence on α . This merits theoretical investigation.



Future work

Theory:

- Prove **monotonicity of $\xi(T)$** for points on the continuum.
- Find theoretical **expectations for the GRU quotient** $\langle \ell_{\max} \rangle / N f_I$, as a function of α , on the continuum. Empirically, we know that there are negative- α and positive- α regimes with different α -dependence.

Experiment:

- Apply more careful **finite-size scaling** to simulation data. (Hallway note: I would be delighted to discuss finite-size scaling with a practitioner.)
- Conduct simulations with **off-lattice quenched positions** (Poisson point process). Lebowitz, Lenci, and Spohn 2000 showed that the point distribution for the Bose gas is not Poisson. Yet, this is a step away from the lattice and toward the true point distribution.
- Conduct simulations with **varying (annealed) point positions** on the continuum. This samples from the true point distribution. Software efficiency (namely, finding which points are near to which) requires a hierarchical partitioning of Λ .
- Develop an algorithm to permit **odd winding numbers**.

Vielen Dank für Ihre Aufmerksamkeit!

Thank you for attending!